Published in:

Journal of Optimization Theory and Applications, Vol. 84, No. 1, pp. 207-234, 1995.

A Linear Programming Based Decomposition Approach in Evaluating Priorities from Pairwise Comparisons and Error Analysis[1]

E. TRIANTAPHYLLOU[2]

Communicated by R. E. Kalaba

---

[1] The author would like to thank Professors Stuart H. Mann, Pennsylvania State University, and Panos M. Pardalos, University of Florida, for their support and valuable comments during the early stages of this research.

[2] Assistant Professor, Department of Industrial and Manufacturing Systems Engineering, Louisiana State University, 3134C CEBA Building, Baton Rouge, Louisiana 70803-6409, U.S.A.
E-mail: IETRIAN@LSUVM.SNCC.LSU.EDU
Web: http://www.imse.lsu.edu/vangelis

Abstract.
One of the most difficult issues in many real life decision- making problems is how to estimate the pertinent data. An approach which uses pairwise comparisons was proposed by Saaty and is widely accepted as an effective way of determining these data. Suppose that two matrices with pairwise comparisons are available. Furthermore, suppose that there is an overlapping of the elements compared in these two matrices. The problem examined in this paper is how to combine the comparisons of the two matrices in order to derive the priorities of the elements considered in both matrices. A simple approach and a linear programming approach are formulated and analyzed in solving this problem. Computational results suggest that the LP approach, under certain conditions, is an effective way for dealing with this problem. The proposed approach is of critical importance because it can also result in a reduction of the total required number of comparisons.

Key Words.
Pairwise comparisons, eigenvectors, analytic hierarchy process, linear programming, fuzzy sets, membership values, artificial intelligence.

1. Introduction

For long time it has been recognized that an exact description of many real physical situations is virtually impossible. This is due to the high degree of imprecision involved in real world situations. Zadeh (Refs. 1 and 2) proposed fuzzy set theory as the means for quantifying the inherent fuzziness that is present in ill-posed problems. Fuzziness is a type of imprecision which may be associated with sets in which there is no sharp transition from membership to nonmembership (Ref. 3). Examples of fuzzy sets are classes of objects characterized by such adjectives as large, small, serious, simple, approximate, etc. (Ref. 3).

A comprehensive description of the importance of fuzzy set theory in engineering and scientific problems is best illustrated in the more than 1,800 references given in Refs. 4 - 12. Currently, an increasingly large number of researchers has been faced by the problem that either their data or their background knowledge is fuzzy. It is particularly critical to people building expert systems and decision support systems, for the knowledge they are dealing with is almost always riddled with vague concepts and judgmental rules (e.g. Refs. 13 - 17). Some recent developments of fuzzy theory on decision making problems are reported in Ref. 18. The most critical step in any application of fuzzy set theory is to effectively estimate the pertinent membership values. Although this is a fundamental problem, there is not a unique way of determining membership values in a fuzzy set. This is mainly due to the way different researchers perceive this problem.

A method proposed by Saaty (Refs. 19 - 21), which is based on pairwise comparisons, has captured the interest of many researchers (see, for example, Refs. 22 - 25). According to this approach the decision maker needs to compare the elements of a fuzzy set by considering all possible pairs. That is, if there are N members then he needs to perform Nx(N-1)/2 pairwise comparisons. Each comparison reflects the personal judgment of the decision maker on how strongly one member of a set belongs to that set when it is compared with another member of the same set.

Saaty (Ref. 20) proposes the use of the scale depicted in Table 1 as the means for quantifying these pairwise comparisons. Other researchers have proposed alternative scales (e.g. Refs. 16 and 27). A scale is nothing but a mapping between a set of discrete linguistic phrases (such as *"A is more important than B"*) and a set of numerical intensities. In Ref. 28 two classes of a total of 78 scales were studied in terms of two evaluative criteria. In that study it was found that no single scale is the best, but different scales are more effective under different circumstances. However, the original Saaty scale is the most widely used and thus this is the one used in the computational experiments described in this paper.

After the pairwise comparisons on N entities are quantified by using a scale, they are used to form the entries of a reciprocal matrix of order N. The (i,j) entry of this matrix reflects the pairwise comparison of the i-th entity when it is compared with the j-th entity. Apparently, entry $a_{i,i} = 1$ and $a_{i,j} = 1/a_{i,j}$. Ideally, if the decision maker is perfectly consistent, then the following relation should also be true:

$$a_{i,j} = a_{i,k} \times a_{j,k}, \quad \text{for any } i,j,k = 1,2,3,\ldots,N.$$

In real life situations there is no guarantee that the above matrix will be consistent. Saaty uses a measurement of consistency, called the Consistency Index (CI) to express the consistency of the comparisons. If the CI coefficient is not within certain acceptable limits, then the

Table 1. Scale of relative importances, according to Saaty (Ref. 21).

Intensity of Importance	Definition	Explanation
1	Equal importance	Two activities contribute equally to the objective
3	Weak importance of one over another	Experience and judgment slightly favor one activity over another
5	Essential or strong importance	Experience and judgment strongly favor one activity over another
7	Demonstrated importance	An activity is strongly favored and its dominance demonstrated in practice
9	Absolute importance	Evidence favoring one activity over another is of the highest possible order of affirmation
2,4,6,8	Intermediate values between the two adjacent judgments	When compromise is needed
Reciprocals of above nonzero	If activity i has one of the above nonzero numbers assigned to it when compared with activity j, then j has the reciprocal value when compared with i.	

comparisons have to be repeated until an acceptable consistency level is reached (Ref. 29). After the above reciprocal matrix has been established, the principal right eigenvector of this matrix is estimated in order to express the relative priorities of the N entities (Ref. 21). That is, according to Saaty (Ref. 21) the entries of this eigenvector reflect the membership values of the members in the set under consideration. An evaluation of the effectiveness of this eigenvector approach can be found in Ref. 30.

Other researchers have proposed the use of a logarithmic regression model (e.g. Refs. 27, 26, 31, and 32) instead of the eigenvector approach. In Ref. 33 a least squares approach is proposed and compared with some other related methods. However, the eigenvector approach is simple and widely used in the literature. Therefore, this is the method used in the computational experiments described in this paper.

The main problem examined in this paper is how to estimate the priorities (i.e., membership values) of N elements when the pairwise comparisons of two subsets with N_1 and N_2 (where $N_1 + N_2 > N$) elements are known. Solving this problem is of critical importance for a number of reasons. When the number of elements to be processed is large, the number of all possible comparisons is very large. For instance, for a collection of 20 elements one needs to perform 190 (=20x19/2) comparisons. Therefore, finding a way for reducing the total number of comparisons is of great practical importance. One may want to partition a large collection of elements into a number of smaller groups. The elements may be clustered into these groups by placing very similar elements in the same group. In this way, the decision maker will evaluate elements which are more homogenous. This has the potential of deriving more accurate comparisons than when one has to compare very different elements.

Another application comes from the area of performing the union operation on the membership values of two sets. If the pairwise comparisons used to derive these membership values are available, then one may want to utilize them in order to determine the membership values of the union of the two sets. This operation can be applicable when performing the union operation in fuzzy data bases.

A similar problem was examined by Harker in Ref. 34. In that approach the decision

maker starts with a minimum set of N (where N is the number of elements) comparisons. Then an expert system-like approach is developed for determining what should be the next comparisons to be made. In this way the decision maker determines the comparisons in a guided manner. The main difference of the problem examined in this paper, is that the pairwise comparisons are assumed to be clustered into two groups of N_1 and N_2 elements each (where $N_1 + N_2 > N$). Therefore, in the present investigation we assume that we have two complete collections of pairwise comparisons. The first collection has $N_1(N_1 - 1)/2$ and the second has $N_2(N_2 - 1)/2$ comparisons.

In this paper two methods are developed for solving the problem of estimating the priorities from two collections of comparisons. The first method is a simple and straightforward approach, while the second is more sophisticated and uses a linear programming (LP) formulation. The LP approach estimates the missing comparisons of the reciprocal matrix by attempting to minimize the CI coefficient of the matrix defined on all the N elements. The two methods are also evaluated in terms of a forward error analysis. The computational results reveal that when the number of common pairwise comparisons is high enough (i.e., $N_1 + N_2$ is significantly greater than N), then the LP approach is the most reliable approach.

2. Problem Description

The main problem examined in this paper is best described in terms of an example. Suppose that there are 5 entities, say A_1, A_2, A_3, A_4, and A_5, for which a decision maker wishes to find their priorities by using pairwise comparisons. These entities may be the alternatives of a problem and the decision maker wishes to find their relative priorities in terms of a single

decision criterion. Furthermore, suppose that the decision maker has available the pairwise comparisons when these 5 entities are grouped into two sub-sets of 4 and 3 members as follows: The first sub-set has the entities $\{A_1, A_2, A_3, A_4\}$ and the second sub-set is $\{A_3, A_4, A_5\}$.

Let the reciprocal matrices with the pairwise comparisons for the previous two sub-sets be as follows:

$$M_1 = \begin{bmatrix} 1.000 & 2.000 & 1.000 & 0.333 \\ 0.500 & 1.000 & 0.500 & 0.200 \\ 1.000 & 2.000 & 1.000 & 0.333 \\ 3.000 & 5.000 & 3.000 & 1.000 \end{bmatrix}, \quad \textit{and} \quad M_2 = \begin{bmatrix} 1.000 & 0.333 & 0.500 \\ 3.000 & 1.000 & 2.000 \\ 2.000 & 0.500 & 1.000 \end{bmatrix}.$$

In this paper it is always assumed that the pairwise comparisons of the same pair of entities is the same regardless of the matrix. This is the reason why the comparison between A_3 and A_4 is the same (equal to 0.333) in both the M_1 and M_2 matrices.

It can be verified that both of the previous matrices are satisfactorily consistent (their CI values are less than 0.10) and thus can be used to derive the relative priorities of the two groups of elements. Saaty in Ref. 19 suggests that an effective way for estimating the principal right eigenvector for this type of matrices is to first calculate the geometric means of each row and then normalize these means. When this procedure is applied to the previous two matrices M_1 and M_2, the following two vectors P_1 and P_2 with the relative priorities are derived, respectively.

$$P_1 = \begin{bmatrix} 0.1855 \\ 0.9710 \\ 0.1855 \\ 0.5318 \end{bmatrix}, \quad \textit{and} \quad P_2 = \begin{bmatrix} 0.1634 \\ 0.5396 \\ 0.2970 \end{bmatrix}.$$

It can be observed that from the first vector the ratio of the relative priorities of the elements A_3 and A_4 is 0.3488 (=0.1855 / 0.5318), while from the second vector the same ratio is 0.3028 (=0.1634 / 0.5396). Therefore, the question we seek to answer in this example is what are the relative priorities when all the 5 elements are considered together?

We can view the previous two matrices as parts of a larger matrix which is defined on the entire 5 elements. When matrices M_1 and M_2 are combined, the following 5x5 matrix M is derived (where "*" indicates undetermined comparison):

$$M = \begin{bmatrix} 1.000 & 2.000 & 1.000 & 0.333 & * \\ 0.500 & 1.000 & 0.500 & 0.200 & * \\ 1.000 & 2.000 & 1.000 & 0.333 & 0.500 \\ 3.000 & 5.000 & 3.000 & 1.000 & 2.000 \\ * & * & 2.000 & 0.500 & 1.000 \end{bmatrix}$$

In other words, only the comparisons A_1/A_5 and A_2/A_5 are missing.

In general, suppose that there are N entities divided into two sub-sets of N_1 and N_2 elements each (where: $N_1 + N_2 > N$). Without loss of generality, suppose that the K (where: $K = (N_1 + N_2) - N$) last elements in the first sub-set correspond, with the same order, to the first K elements in the second sub-set (as was the case in the previous numerical example with $K = 2 = 4+3 - 5$).

When we consider the matrix with all possible comparisons for the N elements, from the above considerations it follows that the decision maker has available the comparisons:

$a_{i,j}$, for $i,j = 1,2,3, \ldots,N_1$, and

$a_{i,j}$, for $i,j = N_2,(N_2+1),(N_2+2), \ldots,N$.

while the comparisons $a_{i,j}$, for $i = (N_1+1), (N_1+2), \ldots, N$, and $j = 1,2,3, \ldots, (N_2-1)$ are undetermined (recall that $a_{j,i} = 1/a_{i,j}$). In other words, the NxN matrix with the Nx(N-1)/2 comparisons can be viewed to be partitioned into sub-matrices as depicted in Figure 1.

In the next section two procedures are developed for estimating the missing comparisons. The first procedure is a simple and straightforward one, while the second procedure attempts an error minimization and is based on an LP formulation.



Fig. 1. Partitioning of the Nx(N-1)/2 pairwise comparisons.

3. Solution Approaches

3.1. Simple Approach

As it was mentioned in the introduction section, perfectly consistent reciprocal matrices with pairwise comparisons satisfy the following relationship:

$$a_{ij} = a_{ik} \times a_{jk}, \quad \text{for } i,j,k = 1,2,3, \ldots, N. \tag{1}$$

From relationship (1) it follows that, in the perfectly consistent case, the missing comparisons (denoted here as X_{ij}) can be determined as follows (see also Figure 1):

$$X_{ij} = a_{ik} / a_{jk}, \qquad \text{for } i = 1,2,3, \ldots, N\text{-}N_2,$$

$$j = (N_1+1),(N_1+2),(N_1+3), \ldots, N, \text{ and}$$

$$k = (N\text{-}N_2+1), \ldots, N_1. \tag{2}$$

In non consistent cases, the previous relationship (2) is not always true. However, the unknown terms X_{ij} are expected to be as close to the products $a_{ik} \times a_{jk}$ as possible. Therefore, it is reasonable to determine the unknown terms X_{ij} as the averages of all possible products. In other words, a simple way is to calculate the terms X_{ij} as follows:

$$X_{ij} = \frac{\sum_{k=N-N_2+1}^{N_1} a_{ik} / a_{jk}}{(N_1 + N_2 - N)},$$

$$\text{for } i = 1,2,3, \ldots,N\text{-}N_2, \text{ and } j = (N_1+1),(N_1+2),(N_1+3), \ldots,N. \tag{3}$$

After these averages are calculated the missing entries of the entire matrix are estimated. Next, the eigenvector approach (or any other pertinent approach) can be applied on the complete matrix and thus the final priorities of the N entities can be estimated.

Although the above averages can be calculated in a straightforward manner, the above approach fails to capture the requirement for the following relationship (4):

$$X_{ij} \approx (a_{i,i'} \times a_{j',j}) \times X_{i',j'}, \qquad \text{for } i,i' = 1,2,3, \ldots,N\text{-}N_2, \text{ and}$$

$$j,j' = (N_1+1),(N_1+2),(N_1+3), \ldots,N. \tag{4}$$

Note that relationship (4) follows directly from the fact that $a_{ij} = 1/a_{ji}$, for any $i,j = 1,2,3, \ldots,N$.

3.2. Linear Programming Approach

The previous approach can be modified and transformed into a more sophisticated procedure. Consider relationship (3). When one wishes to incorporate relationship (4), then relationship (3) may not hold as an equality but instead it will be satisfied as follows:

$$X_{ij} \approx \frac{\sum_{k=N-N_2+1}^{N_1} a_{ik} / a_{jk}}{(N_1 + N_2 - N)},$$

$$\text{for } i = 1,2,3, \ldots,N\text{-}N_2, \text{ and } j = (N_1+1),(N_1+2),(N_1+3), \ldots,N. \quad (5)$$

That is, now the left-hand-side is approximately equal to the right-hand-side. If we wish to make relationship (5) to be an equality, then an error term, denoted as e_{ij}, can be introduced as follows:

$$X_{ij} = \frac{\sum_{k=N-N_2+1}^{N_1} a_{ik} / a_{jk}}{(N_1 + N_2 - N)} + e_{ij},$$

$$\text{for } i = 1,2,3, \ldots,N\text{-}N_2, \text{ and } j = (N_1+1),(N_1+2),(N_1+3), \ldots,N. \quad (6)$$

Similarly, relationship (4) can be transformed into an equality by introducing an error term, denoted as $e_{ij}^{i'j'}$, as follows:

$$X_{ij} = (a_{i,i'} \times a_{j',j}) \times X_{i'j'} + e_{ij}^{i'j'}, \quad \text{for } i,i' = 1,2,3, \ldots,N\text{-}N_2, \text{ and}$$

$$j,j' = (N_1+1),(N_1+2),(N_1+3), \ldots,N. \quad (7)$$

Relationships (6) and (7) suggest that a reasonable treatment to the problem of estimating the missing entries (and thus determining the final priorities) is to attempt to minimize the sum of all the previous error terms. This is in accordance with the implicit assumption that the decision maker attempts to be as consistent in his judgements as possible. Since the errors may be positive

or negative, we would like to minimize the sum of their absolute values. Thus, this consideration leads to the following LP formulation:

$$\textbf{Min } f = \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} |e_{ij}| + \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} \sum_{i'=1}^{N-N_2} \sum_{j'=N_1+1}^{N} |e_{ij}^{i'j'}|$$

$$s.t. \quad X_{ij} = \frac{\sum_{k=N-N_2+1}^{N_1} a_{ik} / a_{jk}}{(N_1 + N_2 - N)} + e_{ij}, \quad \text{for } i = 1,2,3, \ldots,N-N_2,$$

$$\text{and } j = (N_1+1),(N_1+2),(N_1+3), \ldots,N.$$

$$X_{ij} = (a_{i,i'} \; x \; a_{j,j'}) \, x \, X_{i',j'} + e_{ij}^{i'j'}, \quad \text{for } i,i' = 1,2,3, \ldots,N-N_2,$$

$$\text{and } j,j' = (N_1+1),(N_1+2),(N_1+3), \ldots,N.$$

where all X_{ij} variables are ≥ 0, and all e_{ij} and $e_{ij}^{i'j'}$ variables are unrestricted.

The absolute values in the previous LP model can be eliminated by introducing the following transformations in the body of constraints:

$e_{ij} = p_{ij} - n_{ij}$, for $i = 1,2,3, \ldots,N-N_2$, $j = (N_1+1),(N_1+2),(N_1+3), \ldots,N$,

$e_{ij}^{i'j'} = P_{ij}^{i'j'} - N_{ij}^{i'j'}$, for $i,i' = 1,2,3, \ldots,N-N_2$,

and $j,j' = (N_1+1),(N_1+2),(N_1+3), \ldots,N$.

(where all the new variables p_{ij}, n_{ij}, $P_{ij}^{i'j'}$, and $N_{ij}^{i'j'}$ are ≥ 0.)

and also modifying the objective function f as follows:

$$f = \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} [p_{ij} + n_{ij}] + \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} \sum_{i'=1}^{N-N_2} \sum_{j'=N_1+1}^{N} [P_{ij}^{i'j'} + N_{ij}^{i'j'}].$$

This is true because if, say the actual term e_{ij}, has to be negative then it can be easily seen (from the linear dependencies in the columns of the new constraints) that the value of the variable p_{ij} will be equal to zero (as a nonbasic variable) while the variable n_{ij} will be greater than zero. In other words, $|e_{ij}| = p_{ij} + n_{ij}$, and $|e_{ij}^{i'j'}| = P_{ij}^{i'j'} + N_{ij}^{i'j'}$.

Therefore, the previous LP model takes the following form:

$$\textbf{Min } f = \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} [p_{ij} + n_{ij}] + \sum_{i=1}^{N-N_2} \sum_{j=N_1+1}^{N} \sum_{i'=1}^{N-N_2} \sum_{j'=N_1+1}^{N} [P_{ij}^{i'j'} + N_{ij}^{i'j'}]$$

$$s.t. \quad X_{ij} = \frac{\sum_{k=N-N_2+1}^{N_1} a_{ik} / a_{jk}}{(N_1 + N_2 - N)} + p_{ij} - n_{ij}, \qquad for\ i = 1,2,3,\ldots,N\text{-}N_2,$$

$$and\ j = (N_1+1),(N_1+2),(N_1+3),\ldots,N.$$

$$X_{ij} = (a_{i,i'} \times a_{j,j'}) \times X_{i'j'} + P_{ij}^{i'j'} - N_{ij}^{i'j'}, \quad for\ i,i' = 1,2,3,\ldots,N\text{-}N_2,$$

$$and\ j,j' = (N_1+1),(N_1+2),(N_1+3),\ldots,N.$$

where all X_{ij}, p_{ij}, n_{ij}, *and* $P_{ij}^{i'j'}$, $N_{ij}^{i'j'}$ *variables are* ≥ 0.

From the above considerations it follows that the proposed LP model uses $(N\text{-}N_1)(N\text{-}N_2)[3 + 2(N\text{-}N_1)(N\text{-}N_2)]$ continuous variables and $(N\text{-}N_1)(N\text{-}N_2)[1 + (N\text{-}N_1)(N\text{-}N_2)]$ constraints. It can also easily be seen that if the input pairwise comparisons (i.e., the ones defined on the two sub-groups of N_1 and N_2 members) are perfectly consistent, then at optimality the value of the objective function of the previous LP problem is equal to zero (i.e., all errors vanish). Moreover,

the optimal solution of the X_{ij} variables is given by relationship (3). The previous concepts and issues are further illustrated in the following numerical example.

3.3. Numerical Example

In this example we use the same data as in the illustrative example described in section 2. Therefore, the pertinent LP model will have 14 variables and 6 constraints (since N = 5, N_1 = 4 and N_2 = 3). Note that:

$$\frac{\frac{a_{13}}{a_{53}} + \frac{a_{14}}{a_{54}}}{2} = \frac{\frac{1.000}{2.000} + \frac{0.333}{0.500}}{2} = 0.583.$$

Similarly:

$$\frac{\frac{a_{23}}{a_{53}} + \frac{a_{24}}{a_{54}}}{2} = 0.325.$$

and $(a_{11} \times a_{55}) = 1.000$, $(a_{12} \times a_{55}) = 2.000$.

Therefore, the constraints of the LP formulation for this numerical example are as follows:

$$X_{15} = 0.583 + p_{15} - n_{15}$$

$$X_{25} = 0.325 + p_{25} - n_{25}$$

$$X_{15} = 2.000\, X_{25} + P_{15}^{25} - N_{15}^{25}$$

$$X_{25} = 0.500\, X_{15} + P_{25}^{15} - N_{25}^{15}$$

$$X_{15} = 1.000\, X_{15} + P_{15}^{15} - N_{15}^{15}$$

$$X_{25} = 1.000\, X_{25} + P_{25}^{25} - N_{25}^{25}$$

Observe that the last two constraints are <u>redundant</u>, since we can directly set $P_{15}^{15} = N_{15}^{15} = P_{25}^{15} = N_{25}^{25} = 0$. In general, we can <u>always</u> set $P_{ij}^{ij} = N_{ij}^{ij} = 0$. This observation suggests that the

LP formulation, defined in the previous sub-section, can have fewer variables and constraints.

From the previous discussions it follows that the LP formulation for this example is:

$$\textbf{Min } f = p_{15} + n_{15} + p_{25} + n_{25} + P_{15}^{25} + N_{15}^{25} + P_{25}^{15} + N_{25}^{15}$$

$$\begin{aligned} s.t. \quad & X_{15} - p_{15} + n_{15} = 0.583 \\ & X_{25} - p_{25} + n_{25} = 0.325 \\ & X_{15} - 2.000\, X_{25} - P_{15}^{25} + N_{15}^{25} = 0 \\ & X_{25} - 0.500\, X_{15} - P_{25}^{15} + N_{25}^{15} = 0 \\ & X_{15}, X_{25}, p_{15}, p_{25}, n_{15}, n_{25}, P_{15}^{25}, P_{25}^{15}, N_{15}^{25}, N_{25}^{15} \geq 0. \end{aligned}$$

An optimal solution to this LP problem is: $X_{15}^{*} = 0.5830$, $X_{25}^{*} = 0.2915$, $n_{25}^{*} = 0.0335$, and all other variables are equal to zero. The value of the objective function at optimality is: 0.0335.

From this optimal solution it follows that the complete 5x5 matrix, with **all** the pairwise comparisons, is as follows:

$$M' = \begin{bmatrix} 1.000 & 2.000 & 1.000 & 0.333 & 0.583 \\ 0.500 & 1.000 & 0.500 & 0.200 & 0.292 \\ 1.000 & 2.000 & 1.000 & 0.333 & 0.500 \\ 3.000 & 5.000 & 3.000 & 1.000 & 2.000 \\ 1.715 & 3.431 & 2.000 & 0.500 & 1.000 \end{bmatrix}$$

When the eigenvector approach is applied on the previous pairwise comparisons, the following priorities are derived for the entire set of the 5 elements A_1, A_2, A_3, A_4, and A_5.

$$P' = \begin{vmatrix} 0.1392 \\ 0.0722 \\ 0.1350 \\ 0.4137 \\ 0.2398 \end{vmatrix}.$$

If the non-LP approach is used, then it can be easily verified that $X_{15} = 0.583$, and $X_{25} = 0.325$. Therefore, M'' and P'', the matrix with the pairwise comparisons and vector with the priorities, respectively, are as follows:

$$M'' = \begin{vmatrix} 1.000 & 2.000 & 1.000 & 0.333 & 0.583 \\ 0.500 & 1.000 & 0.500 & 0.200 & 0.325 \\ 1.000 & 2.000 & 1.000 & 0.333 & 0.500 \\ 3.000 & 5.000 & 3.000 & 1.000 & 2.000 \\ 1.715 & 3.077 & 2.000 & 0.500 & 1.000 \end{vmatrix},$$

and $$P'' = \begin{vmatrix} 0.1478 \\ 0.0783 \\ 0.1433 \\ 0.4390 \\ 0.1916 \end{vmatrix}.$$

4. Computational Experiments

One challenging issue in designing computational experiments is how to generate the pertinent data. In this study we follow a similar strategy, in deriving the pertinent data, as in the experiments reported in Refs. 33, 35, 36, 37, 38, and 39.

The following forward error analysis is based on the assumption that in the real world the actual relative priorities of the members of collection of entities take on continuous values. These relative priorities can also be viewed as the degree of belonging to a fuzzy set. This continuity assumption is believed to be a reasonable one since it captures the majority of real world cases. As it was mentioned earlier, these members may be a set of alternatives and the

membership values are the degrees that these alternatives meet a single decision criterion.

Let $\omega_1,\omega_3,\omega_3, \ldots, \omega_n$ be the real (and thus unknown) membership values of a fuzzy set with n members. If the decision maker knew the above real values, then he would be able to have constructed a matrix with the real pairwise comparisons. In this matrix, say matrix A, the entries are $\alpha_{ij} = \omega_i / \omega_j$. This matrix is called the Real Continuous Pairwise matrix, or the RCP matrix. Since in the real world the ω_i's are unknown, so are the entries α_{ij} of the previous matrix. However, we can assume here that the decision maker instead of an unknown entry α is able to determine the closest values taken from the set: {1/9,1/8,…,1,2,…,8,9} (if the original Saaty scale is to be used). That is, instead of the real (and thus unknown) value α one is able to determine a_{ij} such that:

the difference $| \alpha - a_{ij} |$ is minimum

and $a_{ij} \in \{1/9,1/8,1/7,\ldots,1,2,\ldots,7,8,9\}$.

In other words, it is assumed here that one's judgments about the value of the pairwise comparison of the i-th element when it is compared with the j-th one, is so accurate that in real life it is the closest (in absolute value terms) of the values one is supposed to choose from.

The matrix with the a_{ij} entries that we assume the decision maker is able to construct has entries from the discrete and finite set: {1/9,1/8,…,1,2,…,8,9}. This second matrix is called the Closest Discrete Pairwise matrix or the CDP matrix. More on some interesting properties of RCP and CDP matrices can be found in Ref. 28.

For illustrative purposes suppose that the actual relative priorities of a set of 5 elements, denoted as $\{A_1, A_2, A_3, A_4, \text{and } A_5\}$, are as follows:

$$\Omega = \begin{vmatrix} 0.1328 \\ 0.0745 \\ 0.1542 \\ 0.3888 \\ 0.2498 \end{vmatrix}$$

Using the previous data, the corresponding RCP matrix can be found to be as follows:

$$RCP = \begin{vmatrix} 1.000 & 1.783 & 0.861 & 0.342 & 0.532 \\ 0.561 & 1.000 & 0.483 & 0.192 & 0.298 \\ 1.161 & 2.070 & 1.000 & 0.397 & 0.617 \\ 2.928 & 5.220 & 2.521 & 1.000 & 1.556 \\ 1.881 & 3.354 & 1.620 & 0.643 & 1.000 \end{vmatrix}$$

This is true because, say the element (1,2), is equal to 1.783 ($= \omega_1 / \omega_2 = 0.1328 / 0.0745$). A similar interpretation holds for the remaining entries in this RCP matrix. Given the previous RCP matrix it can be easily verified that the corresponding CDP matrix is:

$$CDP = \begin{vmatrix} 1.000 & 2.000 & 1.000 & 0.333 & 0.500 \\ 0.500 & 1.000 & 0.500 & 0.200 & 0.333 \\ 1.000 & 2.000 & 1.000 & 0.333 & 0.500 \\ 3.000 & 5.000 & 3.000 & 1.000 & 2.000 \\ 2.000 & 3.000 & 2.000 & 0.500 & 1.000 \end{vmatrix}$$

This is true because, say the element (1,2), is equal to 2.000, since the value 2.000 (taken from the current scale on use) is the closest value to the corresponding entry in the RCP matrix (i.e., 1.783). A similar interpretation holds for the remaining entries in this CDP matrix. Given the previous CDP matrix the relative priorities derived by using the eigenvector approach are:

$$P = \begin{bmatrix} 0.1352 \\ 0.0743 \\ 0.1352 \\ 0.4143 \\ 0.2410 \end{bmatrix}$$

Next, consider the case in which $N_1 = 4$ and $N_2 = 3$. This setting, along with the previous CDP matrix, creates the data considered in the numerical example discussed in section 3.3. In that example it was found that the LP approach and the non-LP approach yield the following relative priorities P_1 and P_2, respectively.

$$P_1 = \begin{bmatrix} 0.1392 \\ 0.0722 \\ 0.1350 \\ 0.4137 \\ 0.2398 \end{bmatrix}, \quad \text{and} \quad P_2 = \begin{bmatrix} 0.1478 \\ 0.0783 \\ 0.1433 \\ 0.4390 \\ 0.1916 \end{bmatrix}.$$

From the previous two sets of relative priorities we can observe that the priorities derived by using the LP approach are closer to the priorities when the CDP matrix is considered (i.e., with vector P). However, the ranking of the 5 elements, when the LP and non-LP approaches are used, is different than the ranking implied by the priorities in P.

The computational experiments were conducted in a similar manner. Sets with $N = 4,5,6, \ldots, 13$ members were assumed. For each case all possible pairs of values of N_1 and N_2 were considered. For instance, for the case with $N = 6$, the (N_1, N_2) pairs were: (3,3), (3,4), (3,5), (4,4), (4,5), and (5,5). In this manner, for the previous values of N a total of 170 different combinations of N, N_1, and N_2 values were generated. These values are depicted in the computational results presented in Table 2 (parts A, B, and C) (note that the results are sorted

in ascending order of the entries in the fifth column).

For each such case, 100 randomly problems were generated and tested to see if the ranking derived by the non-LP and the LP approaches were identical with the ranking derived when the CDP matrix is processed with the eigenvalue method. Note that sometimes the LP formulation with $N > 13$ required excessive CPU time (when 100 random test problems had to be solved), and thus values of $N > 13$ were dropped from further consideration. For each such random problem actual relative weights were assumed as in the numerical example. However, because the Saaty matrices use values from the set: {1/9,1/8,…,1,2,…,8,9} only the random problems which are associated with RCP matrices with entries within the continuous interval [1/9, 9/1] were considered. These computational results are described in more detail in the next section.

5. Computational Results

The computational results are presented in Table 2 (parts A, B, and C). The first three columns in that table present the values of N, N_1, and N_2, respectively. The forth column gives the percent of common pairwise comparisons (PCs) of the number of available comparisons. Furthermore, the fifth column presents the percent of available pairwise comparisons of the total number of comparisons. Finally, the last two columns present the number of contradictions (i.e., when the derived ranking is different than the ranking derived by the original CDP matrix) under the LP and the non-LP approaches.

For instance, consider the seventh row in Table 2 (part A). This row has the numbers: *[3, 5, 6, 8.33, 80, 8, 10]*. In this case the size of the original set is 6. The first sub-set includes the first 3 elements, and the second group includes the last 5 elements. From the values of N, N_1, and N_2 we can verify that the number of available comparisons is 12, the number of common comparisons is 1, and the number of all possible comparisons is 15. Therefore, from the available comparisons, 8.33% are common, and the available comparisons represent 80.00% of all possible

comparisons. When the LP approach was applied on 100 randomly generated test problems with the previous characteristics, in 8 cases the derived ranking was different that the one implied by the corresponding CDP matrices. For the case of using the non-LP approach, however, the same rate was 10%.

These results are also plotted in Figures 2 through 7. Figures 2 and 3 depict the error rates, for sets of different size, when the LP and non-LP approaches are used, respectively. In these figures the error rates are presented as a function of the size of the set (i.e., the N value) and the percent of available comparisons (i.e., the fifth column in Table 2). Figures 4 and 5 also depict the error rates for different sets, but now the percent of common comparisons (i.e., the fourth column in Table 2) are used instead. Figures 6 and 7 present the error rates of the two approaches when the results are in terms of the averages of all set sizes. This is the reason why there are only two curves in figures 6 and 7; one represents the performance of the LP approach, while the other the performance of the non-LP approach. Figure 6 refers to available comparisons, while figure 7 refers to common comparisons.

Table 2a. Computational results, Part A.

N_1	N_2	N	Common PCs (%)	Available PCs (%)	Error rate, LP approach	Error rate, non-LP approach
3	3	4	20	83.33	2	2
3	3	5	0	60	15	5
3	3	6	0	40	82	100
3	4	5	12.5	80	7	5
3	4	6	0	60	21	11
3	4	7	0	42.86	88	100
3	5	6	8.33	80	8	10
3	5	7	0	61.9	27	21
3	5	8	0	46.43	94	100
3	6	7	5.88	80.95	5	8
3	6	8	0	64.29	38	27
3	6	9	0	50	92	100
3	7	8	4.35	82.14	21	19
3	7	9	0	66.67	40	25
3	7	10	0	53.33	98	100
3	8	9	3.33	83.33	15	16
3	8	10	0	68.89	48	41
3	8	11	0	56.36	75	45
3	9	10	2.63	84.44	23	18
3	9	11	0	70.91	57	32
3	9	12	0	59.09	99	100
3	10	11	2.13	85.45	34	27
3	10	12	0	72.73	53	42
3	10	13	0	61.54	99	100
3	11	12	1.75	86.36	33	30
3	11	13	0	74.36	74	48
3	12	13	1.47	87.18	33	35
4	4	5	33.33	90	0	0
4	4	6	9.09	73.33	12	9
4	4	7	0	57.14	21	9
4	4	8	0	42.86	93	100
4	5	6	23.08	86.67	0	2
4	5	7	6.67	71.43	14	15
4	5	8	0	57.14	33	18
4	5	9	0	44.44	94	100
4	6	7	16.67	85.71	3	10
4	6	8	5	71.43	28	18
4	6	9	0	58.33	47	33
4	6	10	0	46.67	100	100
4	7	8	12.5	85.71	7	5
4	7	9	3.85	72.22	32	21
4	7	10	0	60	58	40
4	7	11	0	49.09	98	100
4	8	9	9.68	86.11	13	16
4	8	10	3.03	73.33	23	25
4	8	11	0	61.82	72	42
4	8	12	0	51.52	100	100
4	9	10	7.69	86.67	5	13
4	9	11	2.44	74.55	47	35
4	9	12	0	63.64	76	49

Table 2b. Computational results, Part B.

4	9	13	0	53.85	100	100
4	10	11	6.25	87.27	18	23
4	10	12	2	75.76	51	42
4	10	13	0	65.38	88	61
4	11	12	5.17	87.88	24	26
4	11	13	1.67	76.92	55	44
4	12	13	4.35	88.46	11	19
5	5	6	42.86	93.33	0	1
5	5	7	17.65	80.95	9	11
5	5	8	5.26	67.86	24	16
5	5	9	0	55.56	56	28
5	5	10	0	44.44	98	100
5	6	7	31.58	90.48	4	2
5	6	8	13.64	78.57	12	18
5	6	9	4.17	66.67	38	28
5	6	10	0	55.56	59	35
5	6	11	0	45.45	100	100
5	7	8	24	89.29	4	4
5	7	9	10.71	77.78	18	16
5	7	10	3.33	66.67	35	32
5	7	11	0	56.36	98	100
5	7	12	0	46.97	89	79
5	8	9	18.75	88.89	6	8
5	8	10	8.57	77.78	26	36
5	8	11	2.7	67.27	53	45
5	8	12	0	57.58	84	62
5	8	13	0	48.72	90	75
5	9	10	15	88.89	11	14
5	9	11	6.98	78.18	24	29
5	9	12	2.22	68.18	57	45
5	9	13	0	58.97	95	66
5	10	11	12.24	89.09	11	24
5	10	12	5.77	78.79	43	27
5	10	13	1.85	69.23	55	49
5	11	12	10.17	89.39	13	18
5	11	13	4.84	79.49	43	50
5	12	13	8.57	89.74	18	21
6	6	7	50	95.24	0	0
6	6	8	25	85.71	3	11
6	6	9	11.11	75	19	30
6	6	10	3.45	64.44	44	34
6	6	11	0	54.55	79	51
6	6	12	0	45.45	99	100
6	7	8	38.46	92.86	2	2
6	7	9	20	83.33	10	15
6	7	10	9.09	73.33	39	31
6	7	11	2.86	63.64	57	42
6	7	12	0	54.55	83	52
6	7	13	0	46.15	98	100
6	8	9	30.3	91.67	1	6
6	8	10	16.22	82.22	20	21
6	8	11	7.5	72.73	32	38
6	8	12	2.38	63.64	65	50
6	8	13	0	55.13	97	66
6	9	10	24.39	91.11	5	10
6	9	11	13.33	81.82	20	30
6	9	12	6.25	72.73	45	43
6	9	13	2	64.1	76	51
6	10	11	20	90.91	10	13
6	10	12	11.11	81.82	20	20

Table 2c. Computational results, Part C.

6	10	13	5.26	73.08	47	54
6	11	12	16.67	90.91	6	8
6	11	13	9.38	82.05	29	35
6	12	13	14.08	91.03	14	24
7	7	8	55.56	96.43	0	1
7	7	9	31.25	88.89	6	8
7	7	10	16.67	80	16	22
7	7	11	7.69	70.91	45	36
7	7	12	2.44	62.12	68	56
7	7	13	0	53.85	93	66
7	8	9	44.12	94.44	2	3
7	8	10	25.64	86.67	14	20
7	8	11	13.95	78.18	26	23
7	8	12	6.52	69.7	46	40
7	8	13	2.08	61.54	77	55
7	9	10	35.71	93.33	3	7
7	9	11	21.28	85.45	14	20
7	9	12	11.76	77.27	30	32
7	9	13	5.56	69.23	66	52
7	10	11	29.41	92.73	8	12
7	10	12	17.86	84.85	16	23
7	10	13	10	76.92	41	48
7	11	12	24.59	92.42	7	19
7	11	13	15.15	84.62	20	24
7	12	13	20.83	92.31	9	9
8	8	9	60	97.22	0	0
8	8	10	36.59	91.11	4	6
8	8	11	21.74	83.64	13	22
8	8	12	12	75.76	38	40
8	8	13	5.66	67.95	51	47
8	9	10	48.84	95.56	4	4
8	9	11	30.61	89.09	12	14
8	9	12	18.52	81.82	19	25
8	9	13	10.34	74.36	46	38
8	10	11	40.38	94.55	3	7
8	10	12	25.86	87.88	14	25
8	10	13	15.87	80.77	23	31
8	11	12	33.87	93.94	7	10
8	11	13	22.06	87.18	10	26
8	12	13	28.77	93.59	8	10
9	9	10	63.64	97.78	1	1
9	9	11	41.18	92.73	7	13
9	9	12	26.32	86.36	17	21
9	9	13	16.13	79.49	24	38
9	10	11	52.83	96.36	1	8
9	10	12	35	90.91	14	23
9	10	13	22.73	84.62	17	26
9	11	12	44.44	95.45	4	15
9	11	13	30	89.74	13	22
9	12	13	37.84	94.87	3	6
10	10	11	66.67	98.18	0	0
10	10	12	45.16	93.94	4	10
10	10	13	30.43	88.46	23	26
10	11	12	56.25	96.97	4	3
10	11	13	38.89	92.31	8	16
10	12	13	48	96.15	3	9
11	11	12	69.23	98.48	1	0
11	11	13	48.65	94.87	6	12
11	12	13	59.21	97.44	3	4
12	12	13	71.43	98.72	1	0



Fig. 2. Error rates under the LP approach for sets of different size as a function of the available comparisons.



Fig. 3. Error rates under the non-LP approach for sets of different size as a function of the available comparisons.



Fig. 4. Error rates under the LP approach for sets of different size as a function of the common comparisons.



Fig. 5. Error rates under the non-LP approach for sets of different size as a function of the common comparisons.



Fig. 6. Error rates for the two approaches as a function of the available comparisons.



Fig. 7.

Error rates for the two approaches as a function of the common comparisons.

From the previous results a number of conclusions can be derived. First of all, as it is natural the percent of common comparisons increases with the number of available comparisons. From figures 2 to 5 we can observe that for larger set sizes the error rates are lower. Furthermore, the LP approach yields, on the average, smaller error rates. The effectiveness of the two approaches becomes more transparent in Figures 6 and 7. We can observe that when the percentages of available or common comparisons are low, then either method is unreliable, since it results in too high error rate. For instance, if the percent of available comparisons is less than 50%, then both methods yield error rates higher than 50% (see also Figure 6).

However, for error rates in lower levels, say less than 30%, the LP approach is always the best method. This is the case when the percent of available comparisons is 80% or higher. This is also the case when we consider the percent of common comparisons. Figure 7 suggests that when the percent of common comparisons is 10% or higher, then the two methods have error rates less than 20%. Moreover, the LP approach consistently outperforms the non-LP approach. As it is anticipated, the performance of the methods converges to being perfect (i.e, the error rates vanish) when the percentages of available or common comparisons approach the upper limit of 100%.

The fact that in these results the LP approach performs better than the non-LP approach as the number of common comparisons increases, is in direct agreement with the way the two approaches were designed. The main difference of the two approaches is that the LP approach utilizes the presence of common comparisons in minimizing the sum of the absolute errors. In the non-LP approach, the issue of the common comparisons is totally ignored. However, it is interesting to observe that when the percent of common comparisons is very high (more than 60%), then the non-LP approach is as good as the LP approach. Moreover, the non-LP approach is extremely simpler (we just need to calculate the means in relations (3)) compared to the more CPU time consuming LP formulation.

6. Concluding Remarks

This paper examined the problem of decomposing a large set of pairwise comparisons into two sub-sets. These two sub-sets may have some common comparisons. These sub-sets may be defined by groups of elements which are more similar within these two sub-sets, than when they are considered all together. Two approaches were developed and tested. The first is a simple approach, while the second one is based on an LP formulation. Both approaches first attempt to estimate the missing comparisons and then to derive the relative priorities and rankings by using all the pairwise comparisons.

The LP formulation attempts to minimize the sum of the absolute errors in the missing comparisons. The simulation results suggest that the LP approach is consistently better than the non-LP approach. Furthermore, the effectiveness of this approach improves with the amount of the common comparisons. It should be also stated here that an interesting issue for future research in this area might be how to decompose a large set of elements to be compared into more than two sub-sets.

Finally, it is important to note that the computational results in this paper are contingent to the way the random data were generated. It is possible that if the random data are generated from other distributions, the results to be different. However, it is highly anticipated that the LP approach to perform better than the non-LP approach, and moreover, its performance to improve with the amount of the input data. Since deriving relative priorities from pairwise comparisons is a popular approach in multi-criteria decision making, finding ways of reducing the amount of required data, is of critical importance.

References

1. ZADEH, L.A., *Fuzzy Sets*, Information and Control, Vol. 8, pp. 338-353, 1965.

2. ZADEH, L.A., *Fuzzy Algorithms*, Information and Control, Vol. 12, pp. 94-102, 1968.

3. BELLMAN, R.E., and ZADEH, L.A., *Decision-Making in a Fuzzy Environment*, Management Science, Vol. 17, pp. 140-164, 1970.

4. CHANG, S.K., *Fuzzy Programs*, Proceedings of the Brooklyn Polytechnical Institute Symposium on Computers and Automata, Brooklyn, New York, New York, Vol. 21, pp. 124-135, 1971.

5. DUBOIS, D. and PRADE, H., *Fuzzy Sets and Systems: Theory and Applications*, Academic Press, New York, New York, 1980.

6. GUPTA, M.M., RAGADE, R.K. and YAGER, R.Y., Editors, *Fuzzy Set Theory and Applications*, North-Holland, New York, New York, 1979.

7. XIE, L.A., and BERDOSIAN, S.D., *The Information in a Fuzzy Set and the Relation Between Shannon and Fuzzy Information*, Proceedings of the 7th Annual Conference on Information Sciences and Systems, Johns Hopkins University, Baltimore, Maryland, pp. 212-233, 1983.

8. ZADEH, L.A., FU, K.S., TANAKA, K., and SHIMURA, M., Editors, *Fuzzy Sets and their Applications to Cognitive and Decision Processes*, Academic Press, New York, New York, 1975.

9. ZADEH, L.A., *Fuzzy Sets as a Basis for a Theory of Possibility*, Fuzzy Sets and Systems, Vol. 1, pp. 3-28, 1978.

10. ZADEH, L.A., *A Theory of Approximate Reasoning*, Machine Intelligence, Edited by J. Hayes, D. Michie, and L. I. Mikulich, Wiley, New York, New York, Vol. 9, pp. 149-194, 1979.

11. ZADEH, L.A., *The Concept of a Linguistic Variable and Its Applications to Approximate*

Reasoning, Parts 1-3, Information Sciences, Vol. 8, pp. 199-249, 1975; Vol. 8, pp. 301-357, 1975; Vol. 9, pp. 43-80, 1976.

12. ZIMMERMANN, H.J., *Fuzzy Set Theory - and Its Applications*, Kluwer Academic Publishers, Amsterdam, Holland, 1985.

13. LEE, C.T.R., *Fuzzy Logic and the Resolution Principle*, 2nd International Joint Conference on Artificial Intelligence, London, England, pp. 560-567, 1971.

14. LEE, S.N., GRIZE, Y.L., and DEHNAD, K., *Quantitative Models for Reasoning Under Uncertainty in Knowledge Based Expert Systems*, International Journal of Intelligent Systems, Vol. 2, pp. 15-38, 1987.

15. PRADE, H., and NEGOITA C.V., Editors, *Fuzzy logic in Knowledge Engineering*, Verlag TUV Rheinland, Berlin, Germany, 1986.

16. RAMSAY, A., *Formal Methods in Artificial Intelligence*, Cambridge University Press, Cambridge, England, 1988.

17. ZADEH, L.A., *Can Expert Systems be Designed Without Using Fuzzy Logic?*, Proceedings of the 17th Annual Conference on Information Sciences and Systems, Johns Hopkins University, Baltimore, Maryland, pp. 23-31, 1983.

18. LOOTSMA, F.A., PARDALOS, P.M., and TRIANTAPHYLLOU, E., Editors, *Fuzzy Decision Making*, to be published by Kluwer Academic Publishers, Amsterdam, Holland, 1994.

19. SAATY, T.L., *A Scaling Method for Priorities in Hierarchical Structures*, Journal of Mathematical Psychology, Vol. 15, pp. 234-281, 1977.

20. SAATY, T.L., *Exploring the Interface Between Hierarchies, Multiple Objects and Fuzzy Sets*, Journal of Fuzzy Sets and Systems, Vol. 1, pp. 57-68, 1978.

21. SAATY, T.L., *The Analytic Hierarchy Process*, McGraw Hill International, New York, New York, 1980.

22. CHU, A.T.W., KALABA, R.E., and SPINGARN, K., *A Comparison of Two Methods for Determining the Weights of Belonging to Fuzzy Sets*, Journal of Optimization Theory and Applications, Vol. 27, pp. 531-538, 1979.

23. FEDEROV, V.V., KUZ'MIN, V.B., and VERESKOV, A.I., *Membership Degrees Determination from Saaty Matrix Totalities*, Approximate Reasoning in Decision Analysis, Edited by M. M. Gupta, and E. Sanchez, North-Holland Publishing Company, Amsterdam, Holland, pp. 23-30, 1982.

24. KHURGIN, J.I., and POLYAKOV, V.V., *Fuzzy Analysis of the Group Concordance of Expert Preferences Defined by Saaty Matrices*, Mathematical Research / Mathematische Forschung, Vol. 30, pp. 111-115, 1986.

25. KHURGIN, J.I., and POLYAKOV, V.V., *Fuzzy Approach to the Analysis of Expert Data*, Mathematical Research / Mathematische Forschung, Vol. 30, pp. 116-124, 1986.

26. LOOTSMA, F.A., *Numerical Scaling of Human Judgment in Pairwise-Comparison Methods for Fuzzy Multi-Criteria Decision Analysis*, Mathematical Models for Decision Support, Springer, Berlin, Germany, Vol. 48, pp. 57-88, 1988.

27. LOOTSMA, F.A., MENSCH, T.C.A., and VOS, F.A., *Multi-Criteria Analysis and Budget Reallocation in Long-Term Research Planning*, European Journal of Operational Research, Vol. 47, pp. 293-305, 1990.

28. TRIANTAPHYLLOU, E., F.A. LOOTSMA, P.M. PARDALOS, and S.H. MANN, *On the Evaluation and Application of Different Scales for Quantifying Pairwise Comparisons in Fuzzy Sets*, to appear in: Journal of Multi-Criteria Decision Analysis, 1994.

UPDATED REFERENCE:

TRIANTAPHYLLOU, E., F.A. LOOTSMA, P.M. PARDALOS, and S.H. MANN, *On the Evaluation and Application of Different Scales for Quantifying Pairwise Comparisons in Fuzzy Sets*, Journal of Multi-Criteria Decision Analysis, Vol. 3, No. 3, pp. 133-155,

1994.

29. VARGAS, L.G., *Reciprocal Matrices With Random Coefficients*, Mathematical Modeling, Vol. 3, pp. 69-81, 1982.

30. TRIANTAPHYLLOU, E, and MANN, S.H., *An Examination of the Effectiveness of Multi-Dimensional Decision-Making Methods: A Decision-Making Paradox*, International Journal of Decision Support Systems, Vol. 5, pp. 303-312, 1989.

31. LOOTSMA, F.A., *The French and the American School in Multi-Criteria Decision Analysis*, Recherche Operationnele / Operations Research, Vol. 24, pp. 263-285, 1990.

32. LOOTSMA, F.A., *Scale Sensitivity and Rank Preservation in a Multiplicative Variant of the AHP and SMART*, Report 91-67, Faculty of Technical Mathematics and Informatics, Delft University of Technology, Delft, The Netherlands, 1991.

33. TRIANTAPHYLLOU, E., PARDALOS, P.M., and MANN, S.H., *A Minimization Approach to Membership Evaluation in Fuzzy Sets and Error Analysis*, Journal of Optimization Theory and Applications, Vol. 66, pp. 275-287, 1990.

34. HARKER, P.T., *Incomplete Pairwise Comparisons in the Analytic Hierarchy Process*, Mathematical Modeling, Vol. 9, pp. 837-848, 1987.

35. TRIANTAPHYLLOU, E., and MANN, S.H., *An Evaluation of the Eigenvalue Approach for Determining the Membership Values in Fuzzy Sets*, Fuzzy Sets and Systems, Elsevier Science Publishers, Vol. 35, pp. 295-301, 1990,

36. TRIANTAPHYLLOU, E., and S.H. MANN, *A Computational Evaluation of the AHP and the Revised AHP When the Eigenvalue Method is Used Under a Continuity Assumption*, Computers and Industrial Engineering, 1993, pending publication.

UPDATED REFERENCE:

TRIANTAPHYLLOU, E., and S.H. MANN, *A Computational Evaluation of the AHP and the Revised AHP When the Eigenvalue Method is Used Under a Continuity Assumption*,

Computers and Industrial Engineering, Vol. 26, No. 3, pp. 609-618, 1994.

37. TRIANTAPHYLLOU, E., P.M. PARDALOS, and S.H. MANN, *The Problem of Determining Membership Values in Fuzzy Sets in Real World Situations*, Operations Research and Artificial Intelligence: The Integration of Problem Solving Strategies, Edited by D.E. Brown and C.C. White III, Kluwer Academic Publishers, Amsterdam, Holland, pp. 197-214, 1990.

38. TRIANTAPHYLLOU, E., and SANCHEZ, A., *Identification of the Critical Criteria When Using the Analytic Hierarchy Process*, Working Paper, Department of Industrial and Manufacturing Systems Engineering, Louisiana State University, Baton Rouge, Louisiana, 70803, 1993.

UPDATED REFERENCE:

TRIANTAPHYLLOU, E., and SANCHEZ, A., *A Sensitivity Analysis Approach for Some Deterministic Multi-Criteria Decision-Making Methods*, Decision Sciences, Vol. 28, No. 1, pp. 151-194, 1997.